

Anti-NLRP2 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P107404


Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse
Dilution	WB 1:500~1:1,000; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 665-820 of human NLRP2 (NP_060322.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-NLRP2 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	NBS1; PAN1; NALP2; PYPAF2; CLR19.9; NLRP2.
Calculated MW	Calculated MW: 121 kDa; Observed MW: 120 kDa
Uniprot ID	Q9NX02
Gene ID	55655
Background	This gene is a member of the nucleotide-binding and leucine-rich repeat receptor (NLR) family, and is predicted to contain an N-terminal pyrin effector domain (PYD), a centrally-located nucleotide-binding and oligomerization domain (NACHT) and C-terminal leucine-rich repeats (LRR). Members of this gene family are thought to be important regulators of immune responses. This gene product interacts with components of the I κ B kinase (IKK) complex, and can regulate both caspase-1 and NF- κ B (nuclear factor kappa-light-chain-enhancer of activated B cells) activity. The pyrin domain is necessary and sufficient for suppression of NF- κ B activity. An allelic variant (rs147585490) has been found that is incapable of blocking the transcriptional activity of NF- κ B. Alternative splicing results in multiple transcript variants encoding different isoforms.

Validation Images



Immunofluorescence analysis of NIH/3T3 cells using NLRP2 Rabbit pAb (P107404) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using NLRP2 Rabbit pAb (P107404) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 90s.