

[KO Validated] Anti-DDIT4 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P107967

Product Information

Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:1,000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-232 of human DDIT4 (NP_061931.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-DDIT4 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Dig2; REDD1; REDD-1; T4.
Calculated MW	Calculated MW: 25 kDa; Observed MW: 32 kDa
Uniprot ID	Q9NX09
Gene ID	54541
Background	Predicted to enable 14-3-3 protein binding activity. Involved in defense response to virus; negative regulation of TOR signaling; and response to hypoxia. Located in cytosol.




Western blot analysis of lysates from wild type (WT) and DDIT4 knockout (KO) HeLa cells, using [KO Validated] DDIT4 Rabbit pAb (P107967) at 1:1,000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (SQ201).

Exposure time: 90s.



Western blot analysis of various lysates using [KO Validated] DDIT4 Rabbit pAb (P107967) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (SQ201).

Exposure time: 180s.