

Anti-PIK3CA Mouse mAb

Purified Recombinant Mouse Monoclonal Antibody


Catalog # M900038

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Mouse
Clonality	Monoclonal
Clone No.	95R97F38
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human PIK3CA
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-PIK3CA Mouse mAb [95R97F38] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	PI3-kinase subunit alpha; PI3K-alpha; PI3Kalpha; PtdIns-3-kinase subunit alpha; Phosphoinositide 3-kinase alpha; Phosphoinositide-3-kinase catalytic alpha polypeptide; Serine/threonine protein kinase PIK3CA; PtdIns-3-kinase subunit p110-alpha; p110alpha.
Calculated MW	Calculated MW: 124 kDa; Observed MW: 124 kDa
Uniprot ID	P42336
Gene ID	5290
Background	Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. A pseudogene of this gene has been defined on chromosome 22. [provided by RefSeq, Apr 2016]
Cellular Location	cytoplasm cytosol intercalated disc lamellipodium perinuclear region of cytoplasm phosphatidylinositol 3-kinase complex phosphatidylinositol 3-kinase complex, class IA phosphatidylinositol 3-kinase complex, class IB plasma membrane



Western Blot - Anti-Phospho-NF-kB p65 (Ser536) Rabbit mAb [82S85K62]

All lanes: M900038 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: SH-SY5Y (Human neuroblastoma epithelial cell) whole cell lysates

Lane 3: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lane 4: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 65 kDa

Observed band size: 65 kDa

Developed using the ECL technique (Cat. No. SQ201).



Western Blot - Anti-PIK3CA Mouse mAb [95R97F38]

All lanes: M900038 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: MCF-7 (human breast adenocarcinoma epithelial cell) whole cell lysates

Lane 5: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates

Lane 6: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP Conjugated (Cat. No. LF101) at 1:5,000 dilution

Predicted band size: 124 kDa

Observed band size: 124 kDa

Developed using the ECL technique (Cat. No. SQ201).



Western Blot - Anti-PIK3CA Mouse mAb [95R97F38]

All lanes: M900038 at 1:1,000 dilution

Lane 1: Whole-cell lysates of 293T cells

Lane 2: Lysates from 293T cells overexpressing truncated PIK3CA (52 kDa)


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP Conjugated (Cat. No. LF101) at 1:5,000 dilution

Predicted band size: 52 kDa

Observed band size: 52 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-PIK3CA Mouse mAb [95R97F38]

Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: M900038 at 1:200 dilution


Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-PIK3CA Mouse mAb [95R97F38]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: M900038 at 1:200 dilution

Secondary antibody: Goat Anti-Mouse IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-PIK3CA Mouse mAb [95R97F38]


Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: M900038 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Mouse (488) (Cat. No. LF107) at 1:1,000 dilution (shown in green) and Goat anti-Rabbit (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).



Western Blot - Anti-Phospho-NF-kB p65 (Ser536) Rabbit mAb [82S85K62]

All lanes: M900038 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HeLa cells treated with 10ng/mL TNF-a for 15min


Lysates/proteins at 10 μ g per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 65 kDa

Observed band size: 65 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-Phospho-NF-kB p65 (Ser536) Rabbit mAb [82S85K62]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: M900038 at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green)

Nuclei were stained with DAPI (shown in blue).