

Anti-BMI1 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P100037

Product Information


Application	WB, IP, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000; IP 0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human Bmi1 (NP_005171.4).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-BMI1 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	PCGF4; RNF51; FLVI2/BMI1; flvi-2/bmi-1; Bmi1.
Calculated MW	Calculated MW: 37 kDa; Observed MW: 43 kDa
Uniprot ID	P35226
Gene ID	648
Background	This gene encodes a ring finger protein that is major component of the polycomb group complex 1 (PRC1). This complex functions through chromatin remodeling as an essential epigenetic repressor of multiple regulatory genes involved in embryonic development and self-renewal in somatic stem cells. This protein also plays a central role in DNA damage repair. This gene is an oncogene and aberrant expression is associated with numerous cancers and is associated with resistance to certain chemotherapies. A pseudogene of this gene is found on chromosome X. Read-through transcription also exists between this gene and the upstream COMM domain containing 3 (COMMD3) gene.



Western blot analysis of various lysates using Bmi1 Rabbit pAb (P100037) at 1:5,000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Kit.
 Exposure time: 10s.



Western blot analysis of various lysates using Bmi1 Rabbit pAb (P100037) at 1:5,000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Kit.
 Exposure time: 90s.



Immunoprecipitation analysis of 200 µg extracts of K-562 cells, using 3 µg Bmi1 antibody (P100037). Western blot was performed from the immunoprecipitate using Bmi1 antibody (P100037) at a dilution of 1:1000.