

## Anti-CBL Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P100141

### Product Information


|             |   |
|-------------|---|
| Application | WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, IP, ELISA   |
| Reactivity  | Human, Mouse, Rat   |
| Dilution    | WB 1:1,000~1:5,000; IHC-P 1:50~1:100; IF 1:100~1:500; IP 0.5µg-4µg antibody for 400µg-600µg extracts of whole cells         |
| Host        | Rabbit  |
| Clonality   | Polyclonal  |
| Isotype     | IgG   |
| Label       | Unconjugated  |
| Immunogen   | Recombinant fusion protein containing a sequence corresponding to amino acids 850-906 of human CBL (NP_005179.2).           |
| Format      | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.                     |
| Storage     | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-CBL Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.                       |

### Protein Information


|               |  |
|---------------|--|
| Synonyms      | CBL2; NSLL; C-CBL; RNF55; FRA11B; CBL.   |
| Calculated MW | Calculated MW: 100 kDa; Observed MW: 120 kDa/110 kDa   |
| Uniprot ID    | P22681   |
| Gene ID       | 867  |
| Background    | This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5' UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause of Noonan syndrome-like disorder. |




Immunohistochemistry analysis of paraffin-embedded Human prostate using CBL Rabbit pAb (P100141) at dilution of 1:200 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse lung using CBL Rabbit pAb (P100141) at dilution of 1:200 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunofluorescence analysis of K-562 cells using CBL Rabbit pAb (P100141) at dilution of 1:300 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using CBL Rabbit pAb (P100141) at dilution of 1:300 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of PC-12 cells using CBL Rabbit pAb (P100141) at dilution of 1:300 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from 293T cells, using CBL Rabbit pAb (P100141) at 1:3,000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.  
 Lysates/proteins: 25µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Kit.  
 Exposure time: 30s.



Western blot analysis of lysates from Mouse thymus, using CBL Rabbit pAb (P100141) at 1:3,000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.  
 Lysates/proteins: 25µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Kit.  
 Exposure time: 1s.



Immunoprecipitation analysis of 600 µg extracts of Mouse thymus cells using 3 µg CBL antibody (P100141). Western blot was performed from the immunoprecipitate using CBL antibody at a dilution of 1:1000.