

# Anti-Cyclin H Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P100308

## Product Information

Application	WB, IP, ELISA
Reactivity	Human
Dilution	WB 1:500~1:2,000; IP 0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-323 of human Cyclin H (NP_001230.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Cyclin H Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Synonyms	CAK; p34; p37; Cych; Cyclin H
Calculated MW	Calculated MW: 38 kDa; Observed MW: 36 kDa/
Uniprot ID	P51946
Gene ID	902
Background	<p>The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. A pseudogene of this gene is found on chromosome 4. Alternate splicing results in multiple transcript variants.</p>



Western blot analysis of various lysates using Cyclin H Rabbit pAb (P100308) at 1:1,000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates / proteins: 25  $\mu$ g per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 90s.



Immunoprecipitation analysis of 200  $\mu$ g extracts of K562 cells using 1  $\mu$ g Cyclin H antibody (P100308). Western blot was performed from the immunoprecipitate using Cyclin H antibody (P100308) at a dilution of 1:1000.