

Anti-14-3-3 alpha/beta Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P100373

Product Information


| | |
|-------------|---|
| Application | WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, IP, ELISA |
| Reactivity | Human, Mouse, Rat |
| Dilution | WB 1:500~1:2,000; IHC-P 1:50~1:100; IF 1:50~1:200; IP 0.5μg-4μg antibody for 200μg-400μg extracts of whole cells |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | A synthetic peptide corresponding to a sequence within amino acids 150-246 of human 14-3-3 alpha/beta (NP_003395.1). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-14-3-3 alpha/beta Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information


| | |
|---------------|--|
| Synonyms | HS1; GW128; YWHAA; KCIP-1; HEL-S-1; 14-3-3 alpha/beta. |
| Calculated MW | Calculated MW: 28 kDa; Observed MW: 28 kDa |
| Uniprot ID | P31946 |
| Gene ID | 7529 |
| Background | This gene encodes a protein belonging to the 14-3-3 family of proteins, members of which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals. The encoded protein has been shown to interact with RAF1 and CDC25 phosphatases, suggesting that it may play a role in linking mitogenic signaling and the cell cycle machinery. Two transcript variants, which encode the same protein, have been identified for this gene. |




Immunohistochemistry analysis of paraffin-embedded Human tonsil using 14-3-3 alpha/beta Rabbit pAb (P100373) at dilution of 1:200 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human appendix using 14-3-3 alpha/beta Rabbit pAb (P100373) at dilution of 1:200 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of HeLa cells using 14-3-3 alpha/beta Rabbit pAb (P100373) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using 14-3-3 alpha/beta Rabbit pAb (P100373) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using 14-3-3 alpha/beta Rabbit pAb (P100373) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.