

## Anti-BMI1 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P101591

### Product Information


Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:500~1:2,000; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 120-326 of human Bmi1 (NP_005171.4).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-BMI1 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information


Synonyms	PCGF4; RNF51; FLVI2/BMI1; flvi-2/bmi-1; Bmi1.
Calculated MW	Calculated MW: 37 kDa; Observed MW: 41 kDa
Uniprot ID	P35226
Gene ID	648
Background	This gene encodes a ring finger protein that is major component of the polycomb group complex 1 (PRC1). This complex functions through chromatin remodeling as an essential epigenetic repressor of multiple regulatory genes involved in embryonic development and self-renewal in somatic stem cells. This protein also plays a central role in DNA damage repair. This gene is an oncogene and aberrant expression is associated with numerous cancers and is associated with resistance to certain chemotherapies. A pseudogene of this gene is found on chromosome X. Read-through transcription also exists between this gene and the upstream COMM domain containing 3 (COMMD3) gene.

## Validation Images

---



Immunofluorescence analysis of HeLa cells using Bmi1 Rabbit pAb (P101591).  
Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue:  
DAPI for nuclear staining.



Western blot analysis of lysates from U-937 cells, using Bmi1 Rabbit pAb (P101591) at  
1:1,000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L)  
(LF102) at 1:10,000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3%  
nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 30s.