

Anti-FOXO1/FOXO3A/FOXO4 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P103645

Product Information


Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000; IHC-P 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic peptide of human FOXO1/FOXO3A/FOXO4 (NP_002006.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-FOXO1/FOXO3A/FOXO4 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	FKH 1; FKHI; FKHR; Forkhead (Drosophila) homolog 1 (rhabdomyosarcoma) ; Forkhead box O1; Forkhead box protein O1; Forkhead box protein O1A; Forkhead in rhabdomyosarcoma; Forkhead, Drosophila, homolog of, in rhabdomyosarcoma; FoxO transcription factor; foxo1; FOXO1_HUMAN; FOXO1A; OTTHUMP00000018301; AF6q21; AF6q21 protein; DKFZp781A0677; FKHR2; FKHL1; FKHL1; FKHL1P2; Forkhead (Drosophila) homolog (rhabdomyosarcoma) like 1; Forkhead box O3; Forkhead box O3A; Forkhead box protein O3; Forkhead box protein O3A; Forkhead Drosophila homolog of in rhabdomyosarcoma like 1; Forkhead homolog (rhabdomyosarcoma) like 1; Forkhead in rhabdomyosarcoma like 1; Forkhead in rhabdomyosarcoma-like 1; FOX O3A; FOXO2; foxo3; FOXO3_HUMAN; FOXO3A; MGC12739; MGC31925.
Calculated MW	Observed MW: 80 kDa
Uniprot ID	Q12778, O43524, P98177
Gene ID	2308, 2309, 4303
Background	The Forkhead family of transcription factors mediates many aspects of physiology, including stress response, metabolism, commitment to apoptosis, and development. The Forkhead box subfamily O (FoxO) proteins have garnered particular interest due to their involvement in the modulation of cancer, metabolism and cardiovascular biology. In mammals, four subfamily members have been identified: FOXO1 (previously also known as FKHR), FOXO3 (also known as FKHL1), FOXO4 (also known as AFX), and FOXO6.




Immunohistochemistry analysis of paraffin-embedded Human placenta using FOXO1/FOXO3A/FOXO4 Rabbit pAb (P103645) at dilution of 1:50 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver using FOXO1/FOXO3A/FOXO4 Rabbit pAb (P103645) at dilution of 1:50 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using FOXO1/FOXO3A/FOXO4 Rabbit pAb (P103645) at dilution of 1:50 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Western blot analysis of various lysates using FOXO1/FOXO3A/FOXO4 Rabbit pAb (P103645) at 1:1,000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Kit.
 Exposure time: 180s.