

[KO Validated] Anti-CDK2 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P103661


Product Information

| | |
|-------------|---|
| Application | WB, IHC-P/IF (Tissue-P), ELISA |
| Reactivity | Human, Mouse, Rat |
| Dilution | WB 1:500~1:2,000; IHC-P 1:50~1:200 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | A synthetic peptide corresponding to a sequence within amino acids 200-298 of human CDK2 (NP_001789.2). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-CDK2 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |


Protein Information

| | |
|---------------|--|
| Synonyms | CDKN2; p33(CDK2). |
| Calculated MW | Calculated MW: 34 kDa; Observed MW: 33 kDa |
| Uniprot ID | P24941 |
| Gene ID | 1017 |
| Background | This gene encodes a member of a family of serine/threonine protein kinases that participate in cell cycle regulation. The encoded protein is the catalytic subunit of the cyclin-dependent protein kinase complex, which regulates progression through the cell cycle. Activity of this protein is especially critical during the G1 to S phase transition. This protein associates with and regulated by other subunits of the complex including cyclin A or E, CDK inhibitor p21Cip1 (CDKN1A), and p27Kip1 (CDKN1B). Alternative splicing results in multiple transcript variants. |

Validation Images



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using [KO Validated] CDK2 Rabbit pAb (P103661) at a dilution of 1:200 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Western blot analysis of lysates from wild type (WT) and CDK2 knockout (KO) 293T cells, using [KO Validated] CDK2 Rabbit pAb (P103661) at 1:3,000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 5s.