

Anti-Bub1 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P104215


Product Information

| | |
|-------------|---|
| Application | WB, ELISA |
| Reactivity | Human, Mouse |
| Dilution | WB 1:500~1:1,000 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human BUB1 (NP_004327.1). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-Bub1 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|---------------|---|
| Synonyms | BUB1A; BUB1L; hBUB1; MCPH30; BUB1. |
| Calculated MW | Calculated MW: 122 kDa; Observed MW: 133 kDa |
| Uniprot ID | O43683 |
| Gene ID | 699 |
| Background | This gene encodes a serine/threonine-protein kinase that play a central role in mitosis. The encoded protein functions in part by phosphorylating members of the mitotic checkpoint complex and activating the spindle checkpoint. This protein also plays a role in inhibiting the activation of the anaphase promoting complex/cyclosome. This protein may also function in the DNA damage response. Mutations in this gene have been associated with aneuploidy and several forms of cancer. Alternate splicing results in multiple transcript variants. |

Validation Images



Western blot analysis of various lysates using BUB1 Rabbit pAb (P104215) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.