

Anti-FADD Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P104410

Product Information


Application	WB, IF (Cell)/ICC, IP, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IF 1:50~1:200; IP 0.5µg-4µg antibody for 400µg-600µg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-205 of mouse FADD (NP_034305.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-FADD Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	GIG3; IMD90; MORT1; FADD.
Calculated MW	Calculated MW: 23 kDa; Observed MW: 28 kDa
Uniprot ID	Q61160
Gene ID	14082
Background	Predicted to enable several functions, including caspase binding activity; death effector domain binding activity; and tumor necrosis factor receptor superfamily binding activity. Involved in several processes, including hematopoietic or lymphoid organ development; negative regulation of activation-induced cell death of T cells; and positive regulation of CD8-positive, alpha-beta cytotoxic T cell extravasation. Acts upstream of or within extrinsic apoptotic signaling pathway in absence of ligand; motor neuron apoptotic process; and regulation of programmed cell death. Predicted to be located in several cellular components, including cell body; cytosol; and membrane raft. Predicted to be part of CD95 death-inducing signaling complex and ripoptosome. Predicted to be active in cytoplasm. Is expressed in several structures, including alimentary system; brain; genitourinary system; hemolymphoid system gland; and liver and biliary system. Human ortholog(s) of this gene implicated in leukemia. Orthologous to human FADD (Fas associated via death domain).




Immunofluorescence analysis of paraffin-embedded mouse bone marrow using FADD Rabbit pAb (P104410) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from Mouse heart, using FADD Rabbit pAb (P104410) at 1:1,000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Kit. Exposure time: 10s.



Western blot analysis of lysates from Jurkat cells, using FADD Rabbit pAb (P104410) at 1:1,000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Kit. Exposure time: 180s.



Immunoprecipitation analysis of 600 µg extracts of Mouse heart cells using 3 µg FADD antibody (P104410). Western blot was performed from the immunoprecipitate using FADD antibody (P104410) at a dilution of 1:1000.