

## Anti-IKK alpha Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P104616

### Product Information


Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, IP, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:5,000; IHC-P 1:50~1:200; IF 1:50~1:200; IP 0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 600-700 of human IKK $\alpha$ (NP_001269.3).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-IKK alpha Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information


Synonyms	BPS2; IKK1; IKKA; IKBKA; TCF16; NFKB1A; IKK-alpha; IKK $\alpha$ .
Calculated MW	Calculated MW: 85 kDa; Observed MW: 85 kDa
Uniprot ID	O15111
Gene ID	1147
Background	This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquitination pathway, thereby activating the transcription factor.




Immunohistochemistry analysis of paraffin-embedded Human tonsil using IKK $\alpha$  Rabbit pAb (P104616) at dilution of 1:100 (40 $\times$  lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat liver using IKK $\alpha$  Rabbit pAb (P104616) at dilution of 1:100 (40 $\times$  lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using IKK $\alpha$  Rabbit pAb (P104616) at dilution of 1:100 (40 $\times$  lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunofluorescence analysis of NIH/3T3 cells using IKK $\alpha$  Rabbit pAb (P104616) at dilution of 1:200 (40 $\times$  lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using IKK $\alpha$  Rabbit pAb (P104616) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using IKK $\alpha$  Rabbit pAb (P104616) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates, using IKK $\alpha$  Rabbit pAb (P104616) at 1:2,000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.  
 Lysates/proteins: 25 $\mu$ g per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Kit.  
 Exposure time: 1s.



Immunoprecipitation analysis of 200  $\mu$ g extracts of 293T cells using 3  $\mu$ g IKK $\alpha$  antibody (P104616). Western blot was performed from the immunoprecipitate using IKK $\alpha$  antibody (P104616) at a dilution of 1:1000.