

## [KO Validated] Anti-BRG1 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P104725

### Product Information


Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, IP, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IHC-P 1:50~1:200; IF 1:50~1:200; IP 0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 680-770 of human BRG1/SMARCA4 (NP_003063.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-BRG1 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information


Synonyms	BRG1; CSS4; SNF2; SWI2; MRD16; RTPS2; BAF190; SNF2L4; SNF2LB; hSNF2b; BAF190A; SNF2-beta; BRG1/SMARCA4.
Calculated MW	Calculated MW: 185 kDa; Observed MW: 220 kDa
Uniprot ID	P51532
Gene ID	6597
Background	The protein encoded by this gene is a member of the SWI/SNF family of proteins and is similar to the brahma protein of <i>Drosophila</i> . Members of this family have helicase and ATPase activities and are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin. In addition, this protein can bind BRCA1, as well as regulate the expression of the tumorigenic protein CD44. Mutations in this gene cause rhabdoid tumor predisposition syndrome type 2. Multiple transcript variants encoding different isoforms have been found for this gene.




Immunohistochemistry analysis of paraffin-embedded Human liver cancer using [KO Validated] BRG1/BRG1/SMARCA4 Rabbit pAb (P104725) at dilution of 1:100 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse lung using [KO Validated] BRG1/BRG1/SMARCA4 Rabbit pAb (P104725) at dilution of 1:100 (40× lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of A549 cells using [KO Validated] BRG1/SMARCA4 Rabbit pAb (P104725). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from wild type (WT) and BRG1/BRG1/SMARCA4 knockout (KO) 293T cells, using [KO Validated] BRG1/SMARCA4 Rabbit pAb (P104725) at 1:1,000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1s.



Immunoprecipitation analysis of 300  $\mu$ g extracts of 293T cells using 3  $\mu$ g BRG1/SMARCA4 antibody (P104725). Western blot was performed from the immunoprecipitate using BRG1/SMARCA4 antibody (P104725) at a dilution of 1:1000.