

# Anti-Integrin beta 1 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P105021

## Product Information


Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-728 of human Integrin-β1/CD29 (NP_002202.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Integrin beta 1 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information


Synonyms	CD29; FNRB; MDF2; VLAB; GPIIA; MSK12; VLA-BETA; Integrin-β1/CD29.
Calculated MW	Calculated MW: 88 kDa; Observed MW: 135 kDa
Uniprot ID	P05556
Gene ID	3688
Background	Integrins are heterodimeric proteins made up of alpha and beta subunits. At least 18 alpha and 8 beta subunits have been described in mammals. Integrin family members are membrane receptors involved in cell adhesion and recognition in a variety of processes including embryogenesis, hemostasis, tissue repair, immune response and metastatic diffusion of tumor cells. This gene encodes a beta subunit. Multiple alternatively spliced transcript variants which encode different protein isoforms have been found for this gene.

## Validation Images

---



Immunofluorescence analysis of NIH-3T3 cells using Integrin-β1/CD29 Rabbit pAb (P105021) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using Integrin-β1/CD29 Rabbit pAb (P105021) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit.

Exposure time: 90s.