

Anti-Placental Alkaline Phosphatase Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P105025


Product Information

Application	WB, IP, ELISA
Reactivity	Human
Dilution	WB 1:500~1:2,000; IP 0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 260-460 of human Placental alkaline phosphatase (PLAP) (NP_001623.3).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Placental Alkaline Phosphatase Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	ALP; IAP; ALPI; PALP; PLAP; PLAP-1; Placental alkaline phosphatase (PLAP).
Calculated MW	Calculated MW: 58 kDa; Observed MW: Refer to figures
Uniprot ID	P05187
Gene ID	250
Background	The protein encoded by this gene is an alkaline phosphatase, a metalloenzyme that catalyzes the hydrolysis of phosphoric acid monoesters. It belongs to a multigene family composed of four alkaline phosphatase isoenzymes. The enzyme functions as a homodimer and has a catalytic site containing one magnesium and two zinc ions, which are required for its enzymatic function. One of the main sources of this enzyme is the liver, and thus, it's one of several indicators of liver injury in different clinical conditions. In pregnant women, this protein is primarily expressed in placental and endometrial tissue, however, strong ectopic expression has been detected in ovarian adenocarcinoma, serous cystadenocarcinoma, and other ovarian cancer cells.

Validation Images



Western blot analysis of lysates from HeLa cells, using Placental alkaline phosphatase (PLAP) Rabbit pAb (P105025) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 1s.



Immunoprecipitation analysis of 200 µg extracts of HeLa cells, using 3 µg Placental alkaline phosphatase (PLAP) antibody (P105025). Western blot was performed from the immunoprecipitate using Placental alkaline phosphatase (PLAP) antibody (P105025) at a dilution of 1:1000.