

[KO Validated] Anti-PTP1B Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P105159

Product Information


| | |
|-------------|---|
| Application | WB, ELISA |
| Reactivity | Human, Rat |
| Dilution | WB 1:500~1:1,000 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | Recombinant fusion protein containing a sequence corresponding to amino acids 281-435 of human PTPN1 (NP_002818.1). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-PTP1B Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information


| | |
|---------------|--|
| Synonyms | PTP1B; PTPN1. |
| Calculated MW | Calculated MW: 50 kDa; Observed MW: 50 kDa |
| Uniprot ID | P18031 |
| Gene ID | 5770 |
| Background | The protein encoded by this gene is the founding member of the protein tyrosine phosphatase (PTP) family, which was isolated and identified based on its enzymatic activity and amino acid sequence. PTPs catalyze the hydrolysis of the phosphate monoesters specifically on tyrosine residues. Members of the PTP family share a highly conserved catalytic motif, which is essential for the catalytic activity. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP has been shown to act as a negative regulator of insulin signaling by dephosphorylating the phosphotyrosine residues of insulin receptor kinase. This PTP was also reported to dephosphorylate epidermal growth factor receptor kinase, as well as JAK2 and TYK2 kinases, which implicated the role of this PTP in cell growth control, and cell response to interferon stimulation. Two transcript variants encoding different isoforms have been found for this gene. |



Western blot analysis of lysates from HeLa cells, using [KO Validated] PTPN1 Rabbit pAb (P105159) at 1:600 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Kit.
 Exposure time: 1s.



Western blot analysis of lysates from C6 cells, using [KO Validated] PTPN1 Rabbit pAb (P105159) at 1:600 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Kit.
 Exposure time: 1s.



Western blot analysis of lysates from wild type(WT) and PTPN1 knockout (KO) HeLa(KO) cells, using [KO Validated] PTPN1 Rabbit pAb (P105159) at 1:400 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Kit.
 Exposure time: 1s.