

Anti-PLCG2 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P105373


Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse
Dilution	WB 1:500~1:2,000; IF 1:20~1:100
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human PLC gamma 2 (PLC gamma 2 (PLCG2)) (NP_002652.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-PLCG2 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	FCAS3; APLAID; PLC-IV; PLC-gamma-2; PLC gamma 2 (PLCG2).
Calculated MW	Calculated MW: 148 kDa; Observed MW: 175 kDa
Uniprot ID	P16885
Gene ID	5336
Background	The protein encoded by this gene is a transmembrane signaling enzyme that catalyzes the conversion of 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate to 1D-myo-inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG) using calcium as a cofactor. IP3 and DAG are second messenger molecules important for transmitting signals from growth factor receptors and immune system receptors across the cell membrane. Mutations in this gene have been found in autoinflammation, antibody deficiency, and immune dysregulation syndrome and familial cold autoinflammatory syndrome 3.

Validation Images



Immunofluorescence analysis of Raw264.7 cells using PLC gamma 2 (PLCG2) Rabbit pAb (P105373) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using PLC gamma 2 (PLCG2) Rabbit pAb (P105373) at 1:1,000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.