

[KO Validated] Anti-CTNNA1 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P106396

Product Information


Application	WB, IHC-P/IF (Tissue-P), IP, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000; IHC-P 1:50~1:200; IP 0.5 μ g-4 μ g antibody for 200 μ g-400 μ g extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 637-906 of human α -Catenin (NP_001894.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-CTNNA1 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	α -Catenin; MDBS2; MDPT2; CAP102.
Calculated MW	Calculated MW: 100 kDa; Observed MW: 115 kDa
Uniprot ID	P35221
Gene ID	1495
Background	This gene encodes a member of the catenin family of proteins that play an important role in cell adhesion process by connecting cadherins located on the plasma membrane to the actin filaments inside the cell. The encoded mechanosensing protein contains three vinculin homology domains and undergoes conformational changes in response to cytoskeletal tension, resulting in the reconfiguration of cadherin-actin filament connections. Certain mutations in this gene cause butterfly-shaped pigment dystrophy.




Immunohistochemistry analysis of paraffin-embedded Rat kidney using [KO Validated] α -Catenin Rabbit pAb (P106396) at dilution of 1:100 (40 \times lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human vermiform appendix using [KO Validated] α -Catenin Rabbit pAb (P106396) at dilution of 1:100 (40 \times lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human stomach using [KO Validated] α -Catenin Rabbit pAb (P106396) at dilution of 1:100 (40 \times lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis using [KO Validated] α -Catenin Rabbit pAb (P106396) at dilution of 1:100 (40 \times lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Western blot analysis of various lysates using [KO Validated] α -Catenin Rabbit pAb (P106396) at 1:1,000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 60s.



Western blot analysis of lysates from wild type (WT) and α -Catenin knockout (KO) 293T cells, using [KO Validated] α -Catenin Rabbit pAb (P106396) at 1:1,000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 10s.



Immunoprecipitation analysis of 200 μ g extracts of 293T cells using 3 μ g α -Catenin antibody (P106396). Western blot was performed from the immunoprecipitate using α -Catenin antibody (P106396) at a dilution of 1:500.