

Anti-Complex III Subunit 5 Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P107290

Product Information


Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IHC-P 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 141-274 of human UQCRFS1 (NP_005994.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Complex III Subunit 5 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	RIP1; RIS1; RISP; UQCR5; MC3DN10; UQCRFS1.
Calculated MW	Calculated MW: 30 kDa; Observed MW: 23 kDa
Uniprot ID	P47985
Gene ID	7386
Background	Predicted to enable oxidoreductase activity. Involved in mitochondrial respiratory chain complex III assembly and respiratory electron transport chain. Located in mitochondrion. Part of mitochondrial respiratory chain complex III and mitochondrial respiratory chain complex IV. Implicated in mitochondrial complex III deficiency.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using UQCRFS1 Rabbit pAb (P107290) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Western blot analysis of various lysates using UQCRFS1 Rabbit pAb (P107290) at 1:500 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit (SQ201).
Exposure time: 3s.