

Anti-Phospho-MEK4 (Ser80) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108202

Product Information

| | |
|-------------|---|
| Application | WB, ELISA |
| Reactivity | Human, Mouse, Rat |
| Dilution | WB 1:500~1:2,000 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | A synthetic phosphorylated peptide around S80 of human MAP2K4 (NP_003001.1). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-Phospho-MEK4 (Ser80) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|---------------|--|
| Synonyms | JNKK; MEK4; MKK4; SEK1; SKK1; JNKK1; SERK1; MAPKK4; PRKMK4; SAPKK1; SAPKK-1; Phospho-MAP2K4-S80. |
| Calculated MW | Calculated MW: 44 kDa; Observed MW: 44 kDa |
| Uniprot ID | P45985 |
| Gene ID | 6416 |
| Background | This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants. |



Western blot analysis of lysates from NIH/3T3 and C6 cells, using Phospho-MAP2K4-S80 Rabbit pAb (P108202) at 1:1,000 dilution. NIH/3T3 cells were treated by PDGF (100ng/mL) for 30 minutes after serum-starvation overnight or treated by PDGF (50 ng/mL) for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.

Detection: ECL Enhanced Kit (SQ201).

Exposure time: 150s.