

Anti-Phospho-PDPK1 (Ser241) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108262

Product Information


Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around S241 of human PDK1/PDPK1 (NP_002604.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-PDPK1 (Ser241) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	PDHK1; PDK1; Pyruvate dehydrogenase kinase isoform 1; PDH kinase 1; PDPK1; PRO0461; Phospho-PDK1/PDPK1-S241.
Calculated MW	Calculated MW: 63 kDa; Observed MW: 60-70 kDa
Uniprot ID	O15530
Gene ID	5170
Background	Enables 3-phosphoinositide-dependent protein kinase activity; phospholipase activator activity; and phospholipase binding activity. Involved in several processes, including cell surface receptor signaling pathway; regulation of protein kinase activity; and regulation of signal transduction. Acts upstream of or within intracellular signal transduction. Located in cell projection; cytosol; and plasma membrane. Implicated in prostate cancer. Biomarker of lung non-small cell carcinoma.




Western blot analysis of lysates from NIH/3T3 cells, using Phospho-PDK1/PDPK1-S241 pAb (P108262) at 1:1,000 dilution. NIH/3T3 cells were treated by PDGF (100 ng/mL) at 37°C for 30 minutes after serum-starvation overnight.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% BSA.
 Detection: ECL Kit (SQ201).
 Exposure time: 60s.



Immunofluorescence analysis of C6 cells using Phospho-PDK1/PDPK1-S241 Rabbit pAb (P108262) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using Phospho-PDK1/PDPK1-S241 Rabbit pAb (P108262) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using Phospho-PDK1/PDPK1-S241 Rabbit pAb (P108262) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.