

Anti-Phospho-p38 (Thr180/Tyr182) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108276

Product Information


Application	WB, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2,000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around T180 & Y182 of human p38 MAPK (NP_620581.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-p38 (Thr180/Tyr182) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPBI; PRKM14; PRKM15; SAPK2A; p38ALPHA; Phospho-p38 MAPK-T180/Y182.
Calculated MW	Calculated MW: 41 kDa; Observed MW: 41 kDa/43 kDa
Uniprot ID	Q16539
Gene ID	1432
Background	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TABI protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.



Western blot analysis of lysates from NIH/3T3 cells using Phospho-p38 MAPK-T180/Y182 Rabbit pAb (P108276) at 1:1,000 dilution incubated overnight at 4°C. NIH/3T3 cells were treated by UV at room temperature for 15-30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Kit (SQ201). Exposure time: 90s.



Western blot analysis of lysates from C6 cells using Phospho-p38 MAPK-T180/Y182 Rabbit pAb (P108276) at 1:1,000 dilution incubated overnight at 4°C. C6 cells were treated by UV at room temperature for 15-30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Kit (SQ201). Exposure time: 180s.



Western blot analysis of lysates from HeLa cells, using Phospho-MAPK14-T180/Y182 pAb (P108276) at 1:2,000 dilution or MAPK14 antibody (P102106). HeLa cells were treated by TNF-α (20 ng/mL) at 37°C for 30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA. Detection: ECL Kit (SQ201). Exposure time: 1s.