

Anti-Phospho-Aurora A (Thr288) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108278

Product Information

Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:2,000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around T288 of human AURKA (NP_003591.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-Aurora A (Thr288) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	AIK; ARK1; AURA; BTAK; STK6; STK7; STK15; PPP1R47; Phospho-AURKA-T288.
Calculated MW	Calculated MW: 46 kDa; Observed MW: 48 kDa
Uniprot ID	O14965
Gene ID	6790
Background	The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript variants encoding the same protein have been found for this gene.



Western blot analysis of various lysates, using Phospho-AURKA-T288 Rabbit pAb (P108278) at 1:1,000 dilution. HeLa were treated by Hydroxyurea (4mM) or treated by Nocodazole (50ng/mL) for 20 hours. HT-29 cells were treated by Hydroxyurea (4 mM) for 20 hours or treated by Nocodazole (100ng/mL) for 16 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.

Detection: ECL Kit (SQ201).

Exposure time: 90s.