

## Anti-Phospho-GSK3 beta (Ser9) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108342

### Product Information


Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:100~1:500; IHC-P 1:50~1:200; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around S9 of human GSK3 $\beta$ (NP_001139628.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-GSK3 beta (Ser9) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information


Synonyms	GSK3B; gsk-3 $\beta$ ; Phospho-GSK3 $\beta$ -S9.
Calculated MW	Calculated MW: 47 kDa; Observed MW: 46 kDa
Uniprot ID	P49841
Gene ID	2932
Background	The protein encoded by this gene is a serine-threonine kinase belonging to the glycogen synthase kinase subfamily. It is a negative regulator of glucose homeostasis and is involved in energy metabolism, inflammation, ER-stress, mitochondrial dysfunction, and apoptotic pathways. Defects in this gene have been associated with Parkinson disease and Alzheimer disease.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using Phospho-GSK3β-S9 Rabbit pAb (P108342) at a dilution of 1:100 (40× lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using Phospho-GSK3β-S9 Rabbit pAb (P108342) at a dilution of 1:100 (40× lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using Phospho-GSK3β-S9 Rabbit pAb (P108342) at a dilution of 1:100 (40× lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of HeLa CA and HeLa cells using Phospho-GSK3β-S9 Rabbit pAb(P108342) at a dilution of 1:100 (40× lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Phospho-GSK3 $\beta$ -S9 Rabbit pAb(P108342) at a dilution of 1:100 (40 $\times$  lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using Phospho-GSK3 $\beta$ -S9 Rabbit pAb(P108342) at a dilution of 1:100 (40 $\times$  lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from C6 cells using Phospho-GSK3 $\beta$ -S9 Rabbit pAb (P108342) at 1:500 dilution. C6 cells were treated by Calyculin A (100 nM) at 37 $^{\circ}$ C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25  $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 10s.