

Anti-Phospho-CBL (Tyr700) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P108801


Product Information

| | |
|-------------|---|
| Application | WB, ELISA |
| Reactivity | Human |
| Dilution | WB 1:500~1:2,000 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Label | Unconjugated |
| Immunogen | A synthetic phosphorylated peptide around Y700 of human CBL (NP_005179.2). |
| Format | Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3. |
| Storage | Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage. |
| Precautions | Anti-Phospho-CBL (Tyr700) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|---------------|--|
| Synonyms | CBL2; NSLL; C-CBL; RNF55; FRA11B; Phospho-CBL-Y700. |
| Calculated MW | Calculated MW: 100 kDa; Observed MW: 120 kDa |
| Uniprot ID | P22681 |
| Gene ID | 867 |
| Background | This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5' UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause of Noonan syndrome-like disorder. |

Validation Images



Western blot analysis of lysates from Jurkat cells, using Phospho-CBL-Y700 Rabbit pAb (P108801) at 1:2,000 dilution or CBL antibody (P100141). Jurkat cells were treated by Pervanadate (1 mM) at 37°C for 30 minutes.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.

Detection: ECL Kit (SQ201).

Exposure time: 5s.