

Anti-Phospho-c-Jun (Ser63) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108864

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1,000; IHC-P 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around S63 of human Jun (NP_002219.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-c-Jun (Ser63) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	AP1; p39; AP-1; cJUN; c-Jun; Phospho-c-Jun-S63.
Calculated MW	Calculated MW: 36 kDa; Observed MW: 44 kDa
Uniprot ID	P05412
Gene ID	3725
Background	This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. This gene is intronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.



Western blot analysis of lysates from 293T cells, using Phospho-c-Jun-S63 Rabbit pAb (P108864) at 1:1,000 dilution. 293T cells were treated by Anisomycin (25 $\mu\text{g}/\text{mL}$) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25 μg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).


Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Phospho-c-Jun-S63 Rabbit pAb (P108864) at dilution of 1:200 (40 \times lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Phospho-c-Jun-S63 Rabbit pAb (P108864) at dilution of 1:200 (40 \times lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Phospho-c-Jun-S63 Rabbit pAb (P108864) at dilution of 1:200 (40 \times lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.