

# Anti-Phospho-STAT4 (Tyr693) Rabbit pAb

Purified Rabbit Polyclonal Antibody


Catalog # P108898

## Product Information


Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:100~1:500; IHC-P 1:50~1:100
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around Y693 of human STAT4 (NP_003142.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-STAT4 (Tyr693) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Synonyms	SLEB11; Phospho-STAT4-Y693.
Calculated MW	Calculated MW: 86 kDa; Observed MW: 81 kDa
Uniprot ID	Q14765
Gene ID	6775
Background	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is essential for mediating responses to IL12 in lymphocytes, and regulating the differentiation of T helper cells. Mutations in this gene may be associated with systemic lupus erythematosus and rheumatoid arthritis. Alternate splicing results in multiple transcript variants that encode the same protein.



Immunohistochemistry analysis of paraffin-embedded Human breast carcinoma using Phospho-STAT4-Y693 Rabbit pAb (P108898). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Western blot analysis of lysates from HeLa cells, using Phospho-STAT4-Y693 Rabbit pAb (P108898) at 1:500 dilution. HeLa cells were treated by Pervanadate (1 nM) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Kit (SQ201).

Exposure time: 30s.