

## Anti-Phospho-MRE11 (Ser676) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P109022

### Product Information


Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:1,000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around S676 of human MRE11 (NP_005582.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-MRE11 (Ser676) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	ATLD; HNGS1; MRE11A; MRE11B; Phospho-MRE11 (Ser676).
Calculated MW	Calculated MW: 81 kDa; Observed MW: 81 kDa
Uniprot ID	P49959
Gene ID	4361
Background	This gene encodes a nuclear protein involved in homologous recombination, telomere length maintenance, and DNA double-strand break repair. By itself, the protein has 3' to 5' exonuclease activity and endonuclease activity. The protein forms a complex with the RAD50 homolog; this complex is required for nonhomologous joining of DNA ends and possesses increased single-stranded DNA endonuclease and 3' to 5' exonuclease activities. In conjunction with a DNA ligase, this protein promotes the joining of noncomplementary ends in vitro using short homologies near the ends of the DNA fragments. This gene has a pseudogene on chromosome 3. Alternative splicing of this gene results in two transcript variants encoding different isoforms.

## Validation Images

---



Western blot analysis of lysates from HeLa cells using Phospho-MRE11 (Ser676) Rabbit pAb (P109022) at 1:800 dilution. HeLa cells were treated by Calyculin A (200 nM) at 37°C for 18 hours after serum-starvation overnight. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Kit (SQ201).  
Exposure time: 1s.