

Anti-Phospho-CBL (Ser669) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015915

Product Information

Application	WB, ELISA
Reactivity	Human, Mouse
Dilution	WB 1:1,000~1:2,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	95S60K95
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-CBL (S669)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-CBL (Ser669) Rabbit mAb [95S60K95] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	CBL2; RNF55; CBL; E3 ubiquitin-protein ligase CBL; Casitas B-lineage lymphoma proto-oncogene; Proto-oncogene c-Cbl; RING finger protein 55; RING-type E3 ubiquitin transferase CBL; Signal transduction protein CBL.
Calculated MW	Calculated MW: 100 kDa; Observed MW: 120 kDa
Uniprot ID	P22681
Gene ID	867
Background	This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5' UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause of Noonan syndrome-like disorder. [provided by RefSeq, Jul 2016]
Cellular Location	Cytoplasm.Cell membrane.Cell projection.Cilium.Golgi apparatus.Colocalizes with FGFR2 in lipid rafts at the cell membrane.



Western Blot - Anti-Phospho-CBL (Ser669) Rabbit mAb [95S60K95]

All lanes: R015915 at 1:1,000 dilution

Lane 1: Ball-1 (Human B lymphocyte acute leukemia cell) whole cell lysates

Lane 2: K562 (Human chronic myeloid leukemia cell) whole cell lysates

Lane 3: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates

Lane 4: SH-SY5Y (Human neuroblastoma epithelial cell) whole cell lysates

Lane 5: Mouse liver whole tissue lysates (LPS stimulated)

Lysates/proteins at 10 μ g per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 100 kDa

Observed band size: 120 kDa

Developed using the ECL technique (Cat. No. SQ201).