

Anti-CRKL Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015932

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	54D60P42
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human CrkL
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-CRKL Rabbit mAb [54D60P42] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Crk-like protein; CRKL.
Calculated MW	Calculated MW: 34 kDa; Observed MW: 39 kDa
Uniprot ID	P46109
Gene ID	1399
Background	This gene encodes a protein kinase containing SH2 and SH3 (src homology) domains which has been shown to activate the RAS and JUN kinase signaling pathways and transform fibroblasts in a RAS-dependent fashion. It is a substrate of the BCR-ABL tyrosine kinase, plays a role in fibroblast transformation by BCR-ABL, and may be oncogenic.[provided by RefSeq, Jan 2009]



Western Blot - Anti-CRKL Rabbit mAb [54D60P42]

All lanes: R015932 at 1:1,000 dilution

Lane 1: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 2: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 3: K562 (Human chronic myeloid leukemia cell) whole cell lysates

Lane 4: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates

Lane 5: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates

Lane 6: Mouse brain whole tissue lysates

Lane 7: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates

Lane 8: Rat brain whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 34 kDa

Observed band size: 39 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-CRKL Rabbit mAb [54D60P42]

Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R015932 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.