

Anti-ULK1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R015942

Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	67M08P00
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human ULK1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-ULK1 Rabbit mAb [67M08P00] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	KIAA0722; ULK1; Serine/threonine-protein kinase ULK1; Autophagy-related protein 1 homolog; Unc-51-like kinase 1; ATG1; hATG1.
Calculated MW	Calculated MW: 113 kDa; Observed MW: 130 kDa
Uniprot ID	O75385
Gene ID	8408
Background	Serine/threonine-protein kinase involved in autophagy in response to starvation (PubMed:18936157, PubMed:21460634, PubMed:21795849, PubMed:25040165)
Cellular Location	Cytoplasm.Cytosol.Preautophagosomal structure.Under starvation conditions, is localized to punctate structures primarily representing the isolation membrane that sequesters a portion of the cytoplasm resulting in the formation of an autophagosome.
Tissue Location	Ubiquitously expressed. Detected in the following adult tissues: skeletal muscle, heart, pancreas, brain, placenta, liver, kidney, and lung.



Western Blot - Anti-ULK1 Rabbit mAb [67M08P00]

All lanes: R015942 at 1:2,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: 293T (Human embryonic kidney cell) whole cell lysates

Lane 5: K562 (Human chronic myeloid leukemia cell) whole cell lysates

Lane 6: Caco2 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates

Lane 7: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates

Lane 8: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lane 9: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 113 kDa

Observed band size: 130 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-ULK1 Rabbit mAb [67M08P00]

Sample: Paraformaldehyde-fixed, paraffin embedded human lung cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R015942 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-ULK1 Rabbit mAb [67M08P00]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R015942 at 1:100 dilution and α -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).