

Anti-Phospho-IKB alpha (Ser32) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R016106

Product Information

Application	WB, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000
Host	Rabbit
Clonality	Monoclonal
Clone No.	99I18M73
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-IKB alpha (S32)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-IKB alpha (Ser32) Rabbit mAb [99I18M73] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	EDAID2; IKBA; MAD-3; NFKBI; IKBA_HUMAN; NFKBIA; I-kappa-B-alpha (IkB-alpha IkappaBalphabet); Major histocompatibility complex enhancer-binding protein MAD3; MAD3.
Calculated MW	Calculated MW: 36 kDa; Observed MW: 36 kDa
Uniprot ID	P25963
Gene ID	4792
Background	This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease. [provided by RefSeq, Aug 2011]
Cellular Location	Cytoplasm Nucleus Shuttles between the nucleus and the cytoplasm by a nuclear localization signal (NLS) and a CRM1-dependent nuclear export.



Western Blot - Anti-Phospho-IKB alpha (Ser32) Rabbit mAb [99I18M73]

All lanes: R016106 at 1:1,000 dilution

Lane 1: Ball-1 (Human B lymphocyte acute leukemia cell) whole cell lysates

Lane 2: THP-1 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates

Lane 4: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 36 kDa

Observed band size: 36 kDa

Developed using the ECL technique (Cat. No. SQ201).