

[KD Validated] Anti-ATIC Mouse mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # M020703

Product Information


Application	WB, FC
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:5,000; FC 1:200~1:2,000
Host	Mouse
Clonality	Monoclonal
Clone No.	39R03C65
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human ATIC
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 12 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	[KD Validated] Anti-ATIC Mouse mAb [39R03C65] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	ATIC; 5-Aminoimidazole-4-Carboxamide Ribonucleotide Formyltransferase/IMP Cyclohydrolase; PURH; IMPCHASE; AICARFT; Phosphoribosylaminoimidazolecarboxamide Formyltransferase/IMP Cyclohydrolase; AICAR Transformylase/Inosine Monophosphate Cyclohydrolase; Bifunctional Purine Biosynthesis Protein ATIC; 5-Aminoimidazole-4-Carboxamide-1-Beta-D-Ribonucleotide Transformylase/Inosinase; AICAR Formyltransferase/IMP Cyclohydrolase Bifunctional Enzyme; Epididymis Secretory Sperm Binding Protein Li 70p; Bifunctional Purine Biosynthesis Protein PURH; AICARFT/IMPCHASE; HEL-S-70p; AICAR.
Calculated MW	Calculated MW: 65 kDa; Observed MW: 65 kDa
Uniprot ID	P31939
Gene ID	471
Background	Bifunctional enzyme that catalyzes 2 steps in purine biosynthesis.




Flow cytometric analysis of ATIC expression in C2C12 cells using ATIC antibody (M020703, 1:2,000). Green, isotype control; red, ATIC.



Western blotting analysis using ATIC antibody (M020703). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with ATIC antibody (M020703, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody (1:20,000) respectively. Image was developed using ECL Substrate Kit.



Western blotting analysis using ATIC antibody (M020703). ATIC expression in wild-type (WT) and ATIC shRNA knockdown (KD) HeLa cells with 20 μ g of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with ATIC antibody (M020703, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody (1:20,000) respectively. Image was developed using ECL Substrate Kit.



Validation of ATIC knockdown using flow cytometry. Wild-type(WT, Blue) and knockdown(KD, Green) HeLa cells were stained with ATIC antibody (M020703, 1:2,000) and analyzed using BD flow cytometer.