Anti-Phospho-Rb (Thr252) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody

Catalog # R011348


Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	45K05L75
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Thr252 of human Rb
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-Rb (Thr252) Rabbit mAb [45K05L75] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	RB_HUMAN; Retinoblastoma-associated protein; RB1; p105-Rb; p110-RB1; pRb (Rb); pp110; RB transcriptional corepressor 1; pRb; OSRC; PPP1R130; Retinoblastoma.
Calculated MW	Calculated MW: 106 kDa; Observed MW: 106 kDa
Uniprot ID	P06400
Gene ID	5925
Background	The protein encoded by this gene is a negative regulator of the cell cycle and was the first tumor suppressor gene found. The encoded protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic sarcoma. [provided by RefSeq, Jul 2008]
Cellular Location	Nucleus Cytoplasm During keratinocyte differentiation, acetylation by KAT2B/PCAF is required for nuclear localization (PubMed:20940255). Localizes to the cytoplasm when hyperphosphorylated (By similarity).
Tissue Location	Expressed in the retina. Expressed in foreskin keratinocytes (at protein level) (PubMed:20940255).


Validation Images



Western Blot - Anti-Phospho-Rb (Thr252) Rabbit mAb [45K05L75]
All lanes: R011348 at 1:1,000 dilution
Lane 1: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
Lane 2: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution
Predicted band size: 106 kDa
Observed band size: 106 kDa
Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-Rb (Thr252) Rabbit mAb [45K05L75]
Sample: Paraformaldehyde-fixed, paraffin embedded rat stomach tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R011348 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-Rb (Thr252) Rabbit mAb [45K05L75]
Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R011348 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.