Anti-HDAC2 Mouse mAb

Purified Recombinant Mouse Monoclonal Antibody

Catalog # M012946


Product Information

Application	WB, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse (Cell), Rat
Dilution	WB 1:1,000~1:2,000; IF 1:100~1:200
Host	Mouse
Clonality	Monoclonal
Clone No.	91L51L04
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human HDAC2
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20℃. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20℃ storage.
Precautions	Anti-HDAC2 Mouse mAb [91L51L04] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	HD2; KDAC2; RPD3; YAF1; D10Wsu179e; Yy1bp; mRPD3; HDAC2_HUMAN; HDAC2; Protein deacylase HDAC2; 3.5.1.98; HDAC2_MOUSE; YY1 transcription factor-binding protein.
Calculated MW	Calculated MW: 55 kDa; Observed MW: 55 kDa
Uniprot ID	Q92769
Gene ID	3066
Background	This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2010]
Cellular Location	Nucleus Cytoplasm
Tissue Location	Widely expressed; lower levels in brain and lung.

Validation Images



Western Blot - Anti-HDAC2 Mouse mAb [91L51L04]
All lanes: M012946 at 1:1,000 dilution
Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates
Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lane 4: MCF-7 (human breast adenocarcinoma epithelial cell) whole cell lysates
Lane 5: 293T (Human embryonic kidney cell) whole cell lysates
Lane 6: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates
Lane 7: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates
Lane 8: Mouse brain whole tissue lysates
Lane 9: Mouse liver whole tissue lysates
Lane 10: PC-12 (Rat adrenal pheochromocytoma epithelial cell)whole cell lysates
Lane 11: Rat brain whole tissue lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Mouse IgG(H+L), HRP Conjugated (Cat. No. LF101) at 1:5,000 dilution
Predicted band size: 55 kDa
Observed band size: 55 kDa
Developed using the ECL technique (Cat. No. SQ201).



Immunofluorescence - Anti-HDAC2 Mouse mAb [91L51L04]
Sample: HeLa cells
The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.
Primary antibody: M012946 at 1:100 dilution and and α-tubulin Rabbit Monoclonal Antibody (Cat. No. LF213) at 1:100 dilution
Secondary antibody: Goat anti-Mouse (488) at 1:1,000 dilution (shown in green) and Goat anti-Rabbit (555) at 1:1,000 dilution (shown in red)
Nuclei were stained with DAPI (shown in blue).