Anti-Lipin 1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody

Catalog # R014398


Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Rat, Human, Mouse
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	77R30G66
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Lipin 1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Lipin 1 Rabbit mAb [77R30G66] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	EC=3.1.3.4, KIAA0188, Lipin-1, Lpin1, LPIN1_HUMAN, PAP1, Phosphatidate phosphatase LPIN1.
Calculated MW	Calculated MW: 99 kDa; Observed MW: 130 kDa
Uniprot ID	Q14693
Gene ID	23175
Background	This gene encodes a magnesium-ion-dependent phosphatidic acid phosphohydrolase enzyme that catalyzes the penultimate step in triglyceride synthesis including the dephosphorylation of phosphatidic acid to yield diacylglycerol. Expression of this gene is required for adipocyte differentiation and it also functions as a nuclear transcriptional coactivator with some peroxisome proliferator-activated receptors to modulate expression of other genes involved in lipid metabolism. Mutations in this gene are associated with metabolic syndrome, type 2 diabetes, acute recurrent rhabdomyolysis, and autosomal recessive acute recurrent myoglobinuria (ARARM). This gene is also a candidate for several human lipodystrophy syndromes. [provided by RefSeq, Mar 2017]
Cellular Location	Nucleus. Cytoplasm > cytosol. Endoplasmic reticulum membrane.
Tissue Location	Abundant in adipose tissue and skeletal muscle. Lower levels in some portions of the digestive tract.


Validation Images




Western Blot - Anti-Lipin 1 Rabbit mAb [77R30G66]
All lanes: R014398 at 1:1,000 dilution
Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates
Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates
Lane 5: T24 (Human bladder cancer epithelial cell) whole cell lysates
Lane 6: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates
Lane 7: U2OS (Human osteosarcoma epithelial cell) whole cell lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution
Predicted band size: 99 kDa
Observed band size: 130 kDa
Developed using the ECL technique (Cat. No. SQ201).




Immunohistochemistry - Anti-Lipin 1 Rabbit mAb [77R30G66]
Sample: Paraformaldehyde-fixed, paraffin embedded rat testis tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R014398 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Lipin 1 Rabbit mAb [77R30G66]
Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R014398 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Lipin 1 Rabbit mAb [77R30G66]
Sample: Paraformaldehyde-fixed, paraffin embedded mouse kidney tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R014398 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Lipin 1 Rabbit mAb [77R30G66]
Sample: HeLa cells
The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.
Primary antibodies: R014398 at 1:100 dilution and α-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution
Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)
Nuclei were stained with DAPI (shown in blue).