Anti-MIB1 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody

Catalog # R014621


Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	93P43I68
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human MIB1 / DIP1
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20℃. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20℃ storage.
Precautions	Anti-MIB1 Rabbit mAb [93P43I68] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	DAPK-interacting protein 1, Dip 1, DIP-1, Dip1, E3 ubiquitin protein ligase MIB 1, E3 ubiquitin protein ligase MIB1, E3 ubiquitin-protein ligase mib1, KIAA1323, LVNC7, MIB, mib1, MIB1_HUMAN, Mind bomb homolog 1, Mindbomb E3 ubiquitin protein ligase 1, Ubiquitin ligase mind bomb, Zinc finger ZZ type with ankyrin repeat domain protein 2, ZZANK2, ZZZ6.
Calculated MW	Calculated MW: 110 kDa; Observed MW: 110 kDa
Uniprot ID	Q86YT6
Gene ID	57534
Background	E3 ubiquitin-protein ligase that mediates ubiquitination of Delta receptors, which act as ligands of Notch proteins. Positively regulates the Delta-mediated Notch signaling by ubiquitinating the intracellular domain of Delta, leading to endocytosis of Delta receptors.
Cellular Location	Cytoplasm. Cell membrane. Localizes to the plasma membrane (By similarity). According to PubMed:15048887, it is mitochondrial, however such localization remains unclear.
Tissue Location	Widely expressed at low level. Expressed at higher level in spinal cord, ovary, whole brain, and all specific brain regions examined.


Validation Images




Immunofluorescence - Anti-MIB1 Rabbit mAb [93P43I68]
Sample: A431 cells
The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.
Primary antibodies: R014621 at 1:100 dilution and α-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution
Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)
Nuclei were stained with DAPI (shown in blue).



Immunohistochemistry - Anti-MIB1 Rabbit mAb [93P43I68]
Sample: Paraformaldehyde-fixed, paraffin embedded mouse ovary tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R014621 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-MIB1 Rabbit mAb [93P43I68]
Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R014621 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Western Blot - Anti-MIB1 Rabbit mAb [93P43I68]
All lanes: R014621 at 1:1,000 dilution
Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates
Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lane 4: T24 (Human bladder cancer epithelial cell) whole cell lysates
Lane 5: C2C12 (Mouse myoblasts epithelial cell) whole cell lysates
Lane 6: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates
Lane 7: Rat brain whole tissue lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution
Predicted band size: 110 kDa
Observed band size: 110 kDa
Developed using the ECL technique (Cat. No. SQ201).