Anti-Phospho-Hsp27 (Ser78) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody

Catalog # R015026


Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	22I81N46
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-Hsp27 (S78)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-Hsp27 (Ser78) Rabbit mAb [22I81N46] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	Heat shock 27kDa protein, 28 kDa heat shock protein, CMT2F, DKFZp586P1322, epididymis secretory protein Li 102, Estrogen regulated 24 kDa protein, Estrogen-regulated 24 kDa protein, Heat shock 25kDa protein 1, Heat shock 27 kDa protein, Heat shock 27kD protein 1, Heat shock 27kDa protein 1, Heat shock 28kDa protein 1, Heat Shock Protein 27, Heat shock protein beta 1, Heat shock protein beta-1, heat shock protein family B (small) member 1, HEL-S-102, HMN2B, HS.76067, Hsp 25, HSP 27, Hsp 28, Hsp B1, Hsp25, HSP27, Hsp28, HspB1, HSPB1_HUMAN, SRP27, Stress responsive protein 27, Stress-responsive protein 27.
Calculated MW	Calculated MW: 23 kDa; Observed MW: 23 kDa
Uniprot ID	P04792
Gene ID	3315
Background	This gene encodes a member of the small heat shock protein (HSP20) family of proteins. In response to environmental stress, the encoded protein translocates from the cytoplasm to the nucleus and functions as a molecular chaperone that promotes the correct folding of other proteins. This protein plays an important role in the differentiation of a wide variety of cell types. Expression of this gene is correlated with poor clinical outcome in multiple human cancers, and the encoded protein may promote cancer cell proliferation and metastasis, while protecting cancer cells from apoptosis. Mutations in this gene have been identified in human patients with Charcot-Marie-Tooth disease and distal hereditary motor neuropathy. [provided by RefSeq, Aug 2017].
Cellular Location	Cytoplasm, Nucleus. Cytoplasm > cytoskeleton > spindle. Cytoplasmic in interphase cells. Colocalizes with mitotic spindles in


Validation Images




Western Blot - Anti-Phospho-Hsp27 (Ser78) Rabbit mAb [22I81N46]
All lanes: R015026 at 1:1,000 dilution
Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates
Lane 2 : HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
Lane 3 : HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lane 4 : SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution
Predicted band size: 23 kDa
Observed band size: 23 kDa
Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-Hsp27 (Ser78) Rabbit mAb [22I81N46]
Sample: Paraformaldehyde-fixed, paraffin embedded rat stomach tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R015026 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-Hsp27 (Ser78) Rabbit mAb [22I81N46]
Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R015026 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Phospho-Hsp27 (Ser78) Rabbit mAb [22I81N46]
Sample: HeLa cells
The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.
Primary antibodies: R015026 at 1:100 dilution and α-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution
Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)
Nuclei were stained with DAPI (shown in blue).