Anti-Phospho-STAT3 (Ser727) Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody

Catalog # R015031


Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	14I11Q01
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-STAT3 (S727)
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-STAT3 (Ser727) Rabbit mAb [14I11Q01] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	1110034C02Rik, Acute Phase Response Factor, Acute-phase response factor, ADMIO, APRF, AW109958, DNA binding protein APRF, FLJ20882, HIES, MGC16063, Signal transducer and activator of transcription 3 (acute phase response factor), Signal transducer and activator of transcription 3, STAT 3, Stat3, STAT3_HUMAN.
Calculated MW	Calculated MW: 88 kDa; Observed MW: 88 kDa
Uniprot ID	P40763
Gene ID	6774
Background	The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Sep 2015]
Cellular Location	Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4. Constitutive


Validation Images




Western Blot - Anti-Phospho-STAT3 (Ser727) Rabbit mAb [14I11Q01]
All lanes: R015031 at 1:1,000 dilution
Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates
Lane 2 : HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates
Lane 3 : HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lane 4 : SW620 (Human colorectal carcinoma epithelial cell) whole cell lysates
Lane 5 : Caco2 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates
Lane 6 : Jurkat (Human T lymphocytic leukemia cell) whole cell lysates
Lane 7 : 293T (Human embryonic kidney cell) whole cell lysates
Lane 8 : SCC-9 (Human tongue squamous carcinoma epithelial cell) whole cell lysates
Lane 9 : C2C12 (Mouse myoblasts epithelial cell) whole cell lysates
Lane 10 : Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates
Lane 11 : PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution
Predicted band size: 88 kDa
Observed band size: 88 kDa
Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-Phospho-STAT3 (Ser727) Rabbit mAb [14I11Q01]
Sample: Paraformaldehyde-fixed, paraffin embedded rat heart tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R015031 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Phospho-STAT3 (Ser727) Rabbit mAb [14I11Q01]
Sample: Paraformaldehyde-fixed, paraffin embedded human hepatoma tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R015031 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-Phospho-STAT3 (Ser727) Rabbit mAb [14I11Q01]
Sample: HeLa cells
The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.
Primary antibodies: R015031 at 1:100 dilution and α-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution
Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)
Nuclei were stained with DAPI (shown in blue).