Anti-Phospho-PPP1R12A/PPP1R12B/PPP1R12C (Thr696) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P108591


Product Information

Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:2000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic phosphorylated peptide around T696 of human PPP1R12A/PPP1R12B/PPP1R12CPPP1R12A (NP_002471.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-PPP1R12A/PPP1R12B/PPP1R12C (Thr696) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	MYPT1 (phospho Thr696) ; MYPT1 (phospho T696) ; Myosin Phosphatase (phospho T696) ; Myosin Phosphatase (phospho Thr696) ; p-MYPT1 (Thr696) ; Myosin Phosphatase; M130; MBS; MGC133042; Myosin phosphatase target subunit 1; Myosin phosphatase targeting subunit 1; MYPT 1; MYPT1; PPP1R12A; Protein phosphatase 1 regulatory inhibitor subunit 12A; Protein phosphatase 1 regulatory subunit 12A; Protein phosphatase myosin binding subunit; MYPT1_HUMAN; Myosin phosphatase-targeting subunit 1; Protein phosphatase myosin-binding subunit; PPP1R12A; MYPT1; PPP1R12B; PPP1R12C.
Calculated MW	Observed MW: 140 kDa
Uniprot ID	O14974, O60237, Q9BZL4
Gene ID	4659, 4660, 54776
Background	Myosin phosphatase target subunit 1, which is also called the myosin-binding subunit of myosin phosphatase, is one of the subunits of myosin phosphatase. Myosin phosphatase regulates the interaction of actin and myosin downstream of the guanosine triphosphatase Rho. The small guanosine triphosphatase Rho is implicated in myosin light chain (MLC) phosphorylation, which results in contraction of smooth muscle and interaction of actin and myosin in nonmuscle cells. The guanosine triphosphate (GTP)-bound, active form of RhoA (GTP.RhoA) specifically interacted with the myosin-binding subunit (MBS) of myosin phosphatase, which regulates the extent of phosphorylation of MLC. Rho-associated kinase (Rho-kinase), which is activated by GTP. RhoA, phosphorylated MBS and consequently inactivated myosin phosphatase. Overexpression of RhoA or activated RhoA in NIH 3T3 cells increased phosphorylation of MBS and MLC. Thus, Rho appears to inhibit myosin phosphatase through the action of Rho-kinase. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2009]

Validation Images



Western blot analysis of lysates from Jurkat cells, using Phospho-PPP1R12A/PPP1R12B/PPP1R12C-T696 Rabbit pAb (P108591) at 1:1,000 dilution.
Jurkat cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% BSA.
Detection: ECL Kit.
Exposure time: 1s.