Anti-Phospho-RPS6 (Ser235) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P108789


Product Information

Application	WB, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:2000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A phospho specific peptide corresponding to residues surrounding S235 of human RPS6
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-RPS6 (Ser235) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	S6; eS6; Phospho-S6 Ribosomal Protein (RPS6)-S235.
Calculated MW	Calculated MW: 29 kDa; Observed MW: 29 kDa
Uniprot ID	P62753
Gene ID	6194
Background	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.


Validation Images



Western blot analysis of lysates from HeLa cells using Phospho-S6 Ribosomal Protein (RPS6)-S235 Rabbit pAb (P108789) at 1:800 dilution. HeLa cells were treated by 20% FBS at 37°C for 90 minutes after serum starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 30s.



Western blot analysis of lysates from NIH/3T3 cells using Phospho-S6 Ribosomal Protein (RPS6)-S235 Rabbit pAb (P108789) at 1:800 dilution. NIH/3T3 cells were treated by 20% FBS at 37°C for 90 minutes after serum starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 30s.



Western blot analysis of lysates from PC-12 cells using Phospho-S6 Ribosomal Protein (RPS6)-S235 Rabbit pAb (P108789) at 1:800 dilution. PC-12 cells were treated by Anisomycin (25 ug/ml) at 37°C for 30 minutes.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 30s.